



Research Week 2023

Nighttime average and dipping blood pressure can differ based on the temporal distribution of ambulatory measurements at nighttime

Leandro C. Brito¹; Nicole P. Bowles¹; Andrew W. McHill^{1,2}; Sean P.M. Rice^{1,5}; Matthew P. Butler^{1,3}; Jose Rueda⁴; Jonathan S. Emens⁴; Steven A. Shea^{1,5}; Saurabh S. Thosar^{1,2,5,6}

1) Oregon Institute of Occupational Health Sciences, Oregon Health & Science University, Portland, Oregon, USA. 2) School of Nursing, Oregon Health & Science University, Portland, Oregon, USA. 3) Department of Behavioral Neuroscience, Oregon Health & Science University, Portland, Oregon, USA. 4) Portland VA Medical Center, Portland, Oregon, USA; 5) OHSU-PSU School of Public Health Oregon Health & Science University, Portland, Oregon, USA. 6) Knight Cardiovascular Institute, Oregon Health & Science University, Portland, Oregon, USA

Keywords

Ambulatory Blood Pressure Monitoring, Cardiovascular System, Blood Pressure

Abstract

Introduction:

Nighttime ambulatory (BP) and dipping% (nighttime/daytime BP of <0.9 ; non-dipping ≥ 0.9) are independent predictors of adverse cardiovascular events. Standard guidelines recommend at least $\geq 20/7$ daytime/nighttime measurements for reliable ambulatory BP monitoring, but newer reports suggest $\geq 8/4$ daytime/nighttime measurements are sufficient. Considering BP oscillates across the night, the temporal distribution of measurements across the night may impact nighttime BP and dipping%. To test whether this new recommendation holds for extreme examples of temporal distribution, we compared average nighttime BP and dipping% when using BP measurements only in the first (1st-half) vs. only in the second half (2nd-half) of the night.

Methods:

Seventeen females and twenty-six males (50 ± 10 years [mean $\pm$ SD]) without cardiovascular disease or severe sleep disorders wore an ambulatory BP monitor for 24 hours, programmed to measure BP every 20 minutes when awake and every 30 minutes during a self-selected 8-hour time-in-bed for nighttime. We compared the nighttime BP averages and the calculated dipping% when using the first four measurements from the 1st-half of the nighttime, from the 2nd-half of the nighttime, and all measurements during the nighttime (All nighttime, AN). Repeated measures ANOVA was conducted with significance set as $p < 0.05$.

Results:

Systolic BP was higher using 1st-half than 2nd-half but similar to AN (111 ± 9 vs. 107 ± 11 vs. 109 ± 9 mmHg, $p < 0.01$), while systolic BP dipping% using 1st-half was lower than 2nd-half and AN (9.7 ± 7.4 vs. 13.0 ± 7.6 vs. 11.4 ± 6.7 %, $p < 0.01$, respectively). Diastolic BP and diastolic dipping% were similar among the 1st-half, 2nd-half, and AN segment (63 ± 6 vs. 63 ± 7 vs. 62 ± 7 mmHg, $p = 0.19$) and (17.2 ± 7.6 vs. 17.2 ± 8.8 vs. 18.4 ± 8.8 %, $p = 0.16$), respectively.

Conclusion:

In adults without cardiovascular disease or severe sleep disorders, nighttime BP and dipping% may depend upon when BP measurements are taken. The minimum threshold of 4 measurements for a reliable nighttime BP readout should be used cautiously. Support: NIH F32-HL131308, R01HL163232, R01HL125893, R35HL155681, Medical Research Foundation, and OHSU OFDIR fellowship.